

AŞILANMAYAN ERİŞKİNLER

DR ASLI HAYKIR SOLAY
DIŞKAPI YBEAH
KLİMİK Aylık Toplantıları

OLGU 1


H.K.

70 yaşında kadın


Kastamonu

Tarımla uğraşıyor


***Travma,
Sol el 1. parmakta açık kırık,
Stabilizasyon, sutureasyon,
AMC 2x1000 mg po***



***Sol koldan başlayıp
çeneye doğru yayılan kasılma,
yutma güçlüğü,
soğuk terleme,
Analjezik.***



Tetanoz aşısı
At kaynaklı IG 3000 IU
Metronidazol 500 mg iv




TA: 130/80 mmHg

Nabız: 101/dk


Solunum sayısı: 22/dk

SO₂: %94


Ateş: 37,5°C



★ Tetanoz ★




Oral STOP
İV hidrasyon
İnsan kaynaklı IG 500 IU
Metronidazol 4x500 mg iv
Ses ve ışık izolasyonu
Diazepam 20 mg bolus → 2mg/sa infüzyon



Bilinç açık.

Bir kez sol bacakta kasılma oldu.


*Konuşmakla ve dokunmakla sol kolda kasılması
mevcut.*



TA: 140/90 mmHg


SS:13/dk

Sat O2: %97




Oliclinomel N4


*Sol kolda kasılma arttı. Sol gastrokinemius
kasında kasılmalar → Diazepam 4mg/sa
Oligüri → Üriner sonda*




Nabız : 100-110/dk



TA: 170/110 mmHg
Kaptopril



Kardiak arrest
3 dk Resusitasiyon
Nazal O₂
Diazepam: 6 mg/sa → 4 mg/sa



Entübe

Nabız: 100-165/dk

TA: 90/50 mmHg

İdrar çıkışında azalma

Diazepam 3 mg/sa



Entübe


Nabız: 100-145/dk

TA: 80/50 mmHg


Furosemid

Diazepam 3 mg/sa, propofol

Glutamin, human albumin




Exitus



OLGU 2

- G. K.
- 70 yaşında kadın
- Ev hanımı

Öksürük, ateş, titreme, bulantı, kusma.
Özgeçmiş: HT, SVO, PTE



Genel durum orta-kötü, bilinç açık.

Ateş: 38,3°C



TA: 110/60 mmHg

Nabız: 80/dk


Solunum sayısı: 16/dk

Her iki akciğer orta zonda ral

Batın distandü




WBC:29800/L
Hg:11,4/dL
Plt:25000/L
AST:984 U/L
ALT: 419 U/L
Üre: 213 mg/dL
Kre:1,77 mg/dL






Enfeksiyon hastalıkları konsultasyonu
Nazofarengeal sürüntü örneđi
Seftriakson 2x1 gr iv
Klaritromisin 2x500 mg po
Oseltamivir 2x75 mg po



Eksitus



Nazofarengeal sürüntü: influenza A (**H1N1**)

Balgam ve kan kültürlerinde üreme olmadı.

Olgu 3

K.Ş.

75 yaşında erkek

Paraplejik

Sıcak su dökülmesi sonucunda batın sol alt
kadranda, perine, sol uyluk proksimalinde 3.
dereceden yanık.

Böbrek fonksiyonlarında bozulma
Yanık ünitesine yatırıldı.



ATEŞ
ÖKSÜRÜK



Akciğer grf: sađ orta zonda dansite artışı

Balgam ve kan kùltürü

Nazofarengeal sürüntü örneđi

İmipenem 4x500 mg iv

Osetamivir 2x75 mg po



Balgam ve kan kültürlerinde üreme olmadı
Nazofarengeal sürüntü örneđi

Ziyaretçi yasađı olan yanık ünitesi



?

?

?

?

?



SAĞLIK ÇALIŞANLARI İNFLUENZA AŞISI YAPTIRMA ORANLARI

- **Dışkapı Yıldırım Beyazıt EAH**

Doktor: 50 (%6)

Doktor dışı sağlık presoneli: 75 (%2,6)

- **Ankara'daki tüm sağlık kuruluşları**

Doktor: 80

Doktor dışı sağlık presoneli:342

SAĞLIKLI DANIŞANLAR

**AŞILAMA FIRSATI
OLABİLİR Mİ?**

Olgu 1

A.S

70 yaşında erkek hasta

Huzurevinde kalmak için onay başvurusu

- Klinik şikayetler
- Fizik muayene
- Laboratuvar testleri

- Anatomik veya fonksiyonel aspleni
- Geç kompleman (C5-9) komponent yetmezlikleri
- Bakımevlerinde yaşayan kişiler
- Yurtta kalan öğrenciler
- Rutin olarak *N.meningitis* suşları ile karşılaşan laboratuvar çalışanları
- Askeri personel
- Meningokokal hastalığın hiperendemik veya epidemik olduğu ülkelerde yaşayan veya o bölgeye (Örn. kuru sezonda -Aralık ile Haziran ayları arasında- “Menenjit kuşağı” olarak adlandırılan sahra altı Afrika gibi) seyahat edecek kişiler (öz. bölgesel popülasyonla temasları uzun sürecekse)

Olgu 2

B.K.

30 yaşında kadın

Hemşire

Somali'ye görevli olarak gidecek



TC Sağlık Bakanlığı

Türkiye Hıdut ve Sahiller
Sağlık Genel Müdürlüğü

Seyahat Sağlığı

ANASAYFA

SEYAHAT BİLGİLERİ

ÜLKELER

MERKEZLERİMİZ

İLETİŞİM

ÜLKELERİN SAĞLIK DURUMLARI



SEYAHATE Mİ ÇIKACAKSINIZ?

Nereye Gidiyorsunuz?
Lütfen Gitmek İstediğiniz
Ülkeyi Seçin

Seçiniz

444 77 34

SEYAHAT SAĞLIĞI DANIŞMA HATTI

SEYAHAT SAĞLIĞI EL KİTABI

2. BASKI



T.C. SAĞLIK BAKANLIĞI
TÜRKİYE HUDUT VE SAHİLLER SAĞLIK GENEL MÜDÜRLÜĞÜ

Doğu Afrika ülkeleri hastalık riskleri ve korunma önerileri

ÜLKELER	ÖNERİLEN AŞILAR VE SITMA KORUYUCU İLAÇLAR					
	SARIHUMMA	SITMA	MENENJİT	TİFO*	HEPATİT A**	POLİO
BURUNDİ	EVET	EVET	HAYIR	EVET	EVET	HAYIR
CİBUTİ	HAYIR	EVET	HAYIR	EVET	EVET	EVET
ERİTRE	HAYIR***	EVET	EVET	EVET	EVET	EVET
ETİYOPYA	EVET	EVET	EVET	EVET	EVET	EVET
KENYA	EVET	EVET	EVET	EVET	EVET	EVET
MALAVİ	HAYIR	EVET	HAYIR	EVET	EVET	HAYIR
MOZAMBİK	HAYIR	EVET	HAYIR	EVET	EVET	HAYIR
RUANDA	HAYIR	EVET	HAYIR	EVET	EVET	HAYIR
SOMALİ	HAYIR***	EVET	HAYIR	EVET	EVET	EVET
TANZANYA	HAYIR***	EVET	HAYIR	EVET	EVET	HAYIR
UGANDA	EVET	EVET	EVET	EVET	EVET	HAYIR



Travelers' Health

Home

Destinations (245) -

Somalia -

Traveler View

Healthy Travel
Packing List

Somalia +

Clinician View

Travel Notices

Zika Travel
Information +

Find a Clinic +

Disease Directory

Resources +

[CDC](#) > [Home](#) > [Destinations \(245\)](#)

Health Information for Travelers to Somalia Traveler View



On This Page

- [Vaccines and Medicines](#)
- [Stay Healthy and Safe](#)
- [Healthy Travel Packing List](#)
- [Travel Health Notices](#)
- [After Your Trip](#)



Clinician View

All travelers

You should be up to date on routine vaccinations while traveling to any destination. Some vaccines may also be required for travel.

[Routine vaccines](#)

Make sure you are up-to-date on routine vaccines before every trip. These vaccines include measles-mumps-rubella (MMR) vaccine, diphtheria-tetanus-pertussis vaccine, varicella (chickenpox) vaccine, polio vaccine, and your yearly flu shot.



Most travelers

Get travel vaccines and medicines because there is a risk of these diseases in the country you are visiting.

[Hepatitis A](#)

CDC recommends this vaccine because you can get hepatitis A through contaminated food or water in Somalia, regardless of where you are eating or staying.



[Malaria](#)

You will need to take prescription medicine before, during, and after your trip to prevent malaria. Your doctor can help you decide which medicine is right for you, and also talk to you about other steps you can take to prevent malaria. See more detailed information about [malaria in Somalia](#).



[Typhoid](#)

You can get typhoid through contaminated food or water in Somalia. CDC recommends this vaccine for most travelers, especially if you are staying with friends or relatives, visiting smaller cities or rural areas, or if you are an adventurous eater.



Some travelers

Ask your doctor what vaccines and medicines you need based on where you are going, how long you are staying, what you will be doing, and if you are traveling from a country other than the US.

Cholera	Cholera is a risk in parts of Somalia. CDC recommends this vaccine for adults who are traveling to an area of active cholera transmission. Most people do not travel to areas of active cholera transmission. Avoiding unsafe food and water can also prevent many cholera infections. Cholera is rare in travelers but can be severe. Certain factors may increase the risk of getting cholera or having severe disease (more information).	
Hepatitis B	You can get hepatitis B through sexual contact, contaminated needles, and blood products, so CDC recommends this vaccine if you might have sex with a new partner, get a tattoo or piercing, or have any medical procedures.	
Rabies	Rabies can be found in dogs, bats, and other mammals in Somalia, so CDC recommends this vaccine for the following groups: • Travelers involved in outdoor and other activities (such as camping, hiking, biking, adventure travel, and caving) that put them at risk for animal bites. • People who will be working with or around animals (such as veterinarians, wildlife professionals, and researchers). • People who are taking long trips or moving to Somalia • Children, because they tend to play with animals, might not report bites, and are more likely to have animal bites on their head and neck.	
Yellow Fever	Health recommendation: CDC does not recommend yellow fever vaccine for most travelers to Somalia. However, you might consider this vaccine if you are staying a long time or will be heavily exposed to mosquitoes.	



Adults
Let's vaccinate

2009



Türk İç Hastalıkları Uzmanlık Derneği
Cemal Nafiz Sok. No:12/6 Çankaya Ankara
Tel : +90 312 441 98 23
Fax : +90 312 441 99 83

[Erişkin Aşı Takip Kartı]



"Aşı karnesi kavramı
yaşam boyu olmalıdır"



Haydi Büyüklere Aşıya



TV1